

Dosage Chart | Children 6 to 110 pounds



Acetaminophen Dosage (Tylenol) For Fever & Pain

Child's Weight	Infants' UNCONCENTRATED Drops	Children's Suspension Liquid	Junior Strength Chewable Tablets	Regular Strength Tablets
More than pounds lbs	160mg/5mL Syringe (use syringe in mL)	160mg/teaspoon Teaspoon (use dosing cup)	160mg each Tablet	325mg each Tablet
6 lbs	1.25 mL*	-	-	
12 lbs	2.5 mL	1/2 teaspoon	-	
15 lbs	3.2 mL	-	-	
18 lbs	3.75 mL	3/4 teaspoon	-	
21 lbs	4.5 mL	-	-	
24 lbs	5 mL	1 teaspoon	1 tablet	
30 lbs	6.4 mL	-	-	
35 lbs	7.5 mL	1 1/2 teaspoons	1 1/2 tablets	
47 lbs	10 mL	2 teaspoons	2 tablets	1 tablet
58 lbs	12.5 mL	2.5 teaspoons	-	-
70 lbs	15 mL	3 teaspoons	3 tablets	1.5 tablets
82 lbs	17.5 mL	3.5 teaspoons	-	-
95 lbs	20 mL	4 teaspoons	4 tablets	2 tablets
110 lbs	-	-	-	2 tablets

Please note:

***Do not administer Tylenol to children less than 6 weeks old before consulting with physician.**

***May give every 4-6 hours. Do not exceed 5 doses of acetaminophen in 24 hours.**

Dosage Chart | Children 6 to 110 pounds

Ibuprofen Dosage (Advil, Motrin) for Fever & Pain



Child's Weight	Infants' Ibuprofen Concentrated Drops 50 mg	Children's Ibuprofen Suspension 100 mg / 5mL	Junior Strength Tablet 100mg each	Regular Strength Tablet 200 mg each
----------------	---	--	-----------------------------------	-------------------------------------

Lbs	Dropper (use dropper in mL)	Teaspoon (use dosing cup)	Tablet	Tablet
11 lbs	1.25 mL	1/2 teaspoon / 2.5 mL	-	-
17 lbs	1.875 mL	3/4 teaspoon / 3.75 mL	-	-
22 lbs	2.5 mL	1 teaspoon / 5 mL	-	-
26 lbs	3.25 mL	1.25 teaspoons / 6.25 mL	-	-
33 lbs	3.75 mL	1.5 teaspoons / 7.5 mL	1.5 tablets	-
44 lbs	5 mL	2 teaspoons / 10 mL	2 tablets	1 tablet
55 lbs	7.5 mL	2.5 teaspoons / 12.5 mL	2.5 tablets	1 tablet
66 lbs	-	3 teaspoons / 15 mL	3 tablets	1.5 tablets
77 lbs	-	3.5 teaspoons / 17.5 mL	3.5 tablets	1.5 tablets
88 lbs	-	4 teaspoons / 20 mL	4 tablets	2 tablets
99 lbs	-	4.5 teaspoons / 23.5 mL	4.5 tablets	2 tablets
110 lbs	-	5 teaspoons / 25 mL	5 tablets	2.5 tablets
120 lbs	-	-	-	3 tablets

Please note: Do not use Ibuprofen for patients under 6 months of age.

May give every 6-8 hours. Do not exceed 4 doses of ibuprofen in 24 hours.